

NATIONAL STANDARD OF THE PEOPLE'S REPUBLIC OF CHINA

中华人民共和国国家标准

GB/T 18204.1-2000

Methods of microbiological examination for air in public places-Determination of aerobic bacterial count 公共场所空气微生物检验方法细菌总数测定

Issued on September 03, 2000

Implemented on January 01, 2001

Foreword

This Standard was formulated in order to carry out *Public Health Management Regulations*, GB 9663~9673-1996, GB 16153-1996 *Hygienic standard for dining room*, strengthen management of public health. The method of this Standard is consistence with monitoring methods of GB 9663~9673-1996, GB 16153-1996.

Method I of this Standard is arbitration method.

This Standard was issued for the first time.

This standard was proposed by Ministry of Health of the People's Republic of China

Draft units of this Standard are Medical Institute of Environmental Health Sciences
Institute of Chinese Preventive Detection, Beijing Health and Epidemic Prevention Station,
Guangdong Health and Epidemic Prevention Station and Tianjin Health and Epidemic
Prevention Station.

Drafting staffs of this standard: Zhou Shuyu, Chen Xiping, Gao Hui, Cai Wen and Zhang Shulan.

Methods of microbiological examination for air in public places-Determination of aerobic bacterial count

1 Scope

This standard specifies the determination method of aerobic bacterial count for air in public places.

This standard applies to the determination of aerobic bacterial count for air in public places.

2 Definitions

This standard uses the following definitions.

2.1 Impacting method

A kind of sampling method that uses the impact-type Microbial Air Monitoring Systems for sampling; through air exhaust dynamic effect, make the air pass through slit or hole to produce high-speed airflow, so that the bacteria-bearing particles suspending in the air strike the nutrient agar medium plates; after incubating for 48 h under 37°C, calculate bacterial colonies count contained for each cubic meter in the air.

2.2 Natural sinking method

It is refers to the sampling determination method for aerobic bacterial count by exposing the nutrient agar plate with diameter of 9cm at sampling points for 5min, after incubating for 48 h under 37°C.

3 Instruments and equipment

- 3.1 High-pressure steam sterilizer:
- 3.2 Dry heat sterilizer.
- 3.3 Constant temperature incubator
- 3.4 Refrigerator
- 3.5 Plate (diameter of 9cm).
- 3.6 General equipment for preparing medium: graduated cylinder, erlenmeyer flasks, pH meter or precise pH paper, etc.
- 3.7 Impact air microbial samplers.



北京文心雕语翻译有限公司

Beijing Lancarver Translation Inc.

完整版本请在线下单

或咨询:

TEL: 400-678-1309

QQ: 19315219

Email: info@lancarver.com

http://www.lancarver.com

对公账户:

单位名称:北京文心雕语翻译有限公司

开户行:中国工商银行北京清河镇支行

账号: 0200 1486 0900 0006 131

支付宝账户: info@lancarver.com

注:付款成功后,请预留电邮,完整版本将在一个工作日内通过电子 PDF 或Word 形式发送至您的预留邮箱,如需索取发票,下单成功后的三个工作日内安排开具并寄出,预祝合作愉快!